

# Construction of Pbs.PGK.PCR1




FIG. 1

# Construction of pIG.E1a.E1b.X




FIG. 2




FIG. 3A

# Construction of pIG.E1a.NEO





FIG. 3B

FIG. 4

Construction of pIG.E1a.E1b



# Construction of pIG.NEO





FIG. 5

# Overview of available adenovirus packaging constructs and assessment of their capacity to transform primary kidney cells



\*average of 5 plates 21 days after transection

FIG. 6

Western blotting analysis of A549 clones transfected with pIG.E1A.NEO and PER clones (HER cells transfected with pIG.E1A.E1B)




FIG. 7



# Southern blot analyses of 293, 911 and PER cell lines




FIG. 8

Transfection efficiency of PER.C3, PER.C5, PER.C6 and 911 cells. Cells were cultured in 6-well plates and transfected (n=2) with 5  $\mu$ g pRSV.lacZ by calcium-phosphate co-precipitation. Forty-eight hours later the cells were stained with X-GAL. The mean percentage of blue cells is shown.




FIG. 9

# Construction of pMLP1.TK





FIG. 10



New recombinant adenoviruses and packaging constructs without sequence overlap



Packaging system based on established cell lines: transfection with E1a and selection with G418 **FIG. 1 B**

# Generation of recombinant adenovirus




FIG. 12




FIG. 13

Replication of Adenovirus




FIG. 14



The potential hairpin conformation of a single-stranded DNA molecule that contains the HP/asp sequences used in these studies. Restriction with the restriction endonucleases *Asp718I* of plasmid pICLHa, containing the annealed oligonucleotide pair HP/asp1 en HP/asp2 will yield a linear double-stranded DNA fragment. In cells in which the required adenovirus genes are present, replication can initiate at the terminus that contains the ITR sequence. During the chain elongation, the one of the strands will be displaced. The terminus of the single-stranded displaced-strand molecule can adopt the conformation depicted above. In this conformation the free 3'-terminus can serve as a primer for the cellular and/or adenovirus DNA polymerase, resulting in conversion of the displaced strand in a double-stranded form.




FIG. 15




FIG. 16




FIG. 17




FIG. 18




FIG. 19

# Cloned adenovirous fragments




FIG. 20

# Adapter plasmid pAd5/L420-HSA




FIG. 2I

Adapter plasmid pAd5/CLIP




FIG. 22



# Generation of recombinant adenoviruses




FIG. 23

# Minimal adenovirus vector pMV/L420H





FIG. 24

# Construction of pWE/AdΔ5'



1. pcr fragment  
left ITR

pBr/Ad.Cla-Bam  
x EcoRI and  $\text{AfIII}$ (partial)

pBr/Ad.Bam-rITR.pac  
x BamHI and PacI

pAd/LTR( $\Delta 5'$ )-BamHI

+ pWE15.Pac x PacI

xPacI, BamHI

+ pWE/AdΔ5'

FIG. 25




FIG. 26A




FIG. 26C





FIG. 26B




Average percentage CPE efficiency: 86 %

FIG. 27

Gene	Insert kb	Average titer 0.8 ±0.7 x 10 <sup>9</sup> pfu/ml
• ceNOS	3.6	
• hTERT	3.5	
• hTERT D712A	3.5	
• lacZ	3.2	
• hCAT1	2.2	
• GLVR2	2.0	
• Luc	1.7	
• SOD3	1.4	
• MAX1	.550	
• hVEGF121	.511	
• hIL3	.434	
• UBC9	.412	
• ANG1-7	.104	

FIG. 28

% wells producing functional virus



Gene	Number of CPE+ wells
ceNOS	19/19
IL3	7/7
lacZ	36/36
Luc	40/40
GFP	48/48

Gene	Number of plaques
ceNOS	9/9
IL3	9/9
lacZ	40/40
Luc	9/9
EGFP	IP
GLVR2	9/9

FIG. 29






FIG. 30




FIG. 3IA




FIG. 3IB




FIG. 3IC







FIG. 32



AS-virus

FIG. 33




FIG. 34A




FIG. 34B




FIG. 34C




FIG. 34D






FIG. 34E




FIG. 34F

10036949.122101




FIG. 34G




FIG. 34H




FIG. 34 I




FIG. 34J

FIG. 34K




FIG. 34K




FIG. 34L






FIG. 34M

Relative amounts of wells with CPE after transfection of PER.C6/E2A cells with pCLIP-LacZ and the adapter plasmid pIPspAdapt2.




FIG. 34N




FIG. 35

# Construction total Adeno cDNA Library (1)




FIG. 36A

# Construction total Adeno cDNA Library (II)




FIG. 36B

EXAMPLE 21 384 WELL PLATE IN PROGRESS

Co-transfections on 384 well plates




FIG. 37A

Co-transfections on 96 well plates  
(control plate)




FIG. 37B

Co-transfections on 384 well plates




FIG. 37C

Co-transfections on 96 well plates  
(control plate)




FIG. 37D

Medium changed 7 days after transfection




FIG. 38A

Medium not changed




FIG. 38B

Propagation 7 days after transfection




FIG. 38C

Cell titration experiment #1




FIG. 39A

Cell titration experiment #2




FIG. 39B

Cell titration experiment #3




FIG. 39C






FIG. 40




FIG. 41




FIG. 42




FIG. 43




FIG. 44